



An Application of Hayashi's Inequality for Differentiable Functions

R. P. AGARWAL

Department of Mathematics, National University of Singapore, 10 Kent Ridge Crescent, Singapore

S. S. DRAGOMIR

Department of Mathematics, Timisoara University, Timisoara, Romania

(Received April 1996; accepted May 1996)

Abstract—In this paper, we offer an application of Hayashi's generalization of Steffensen's inequality for differentiable functions, as well as its connections with the celebrated Hermite-Hadamard's integral inequality for convex functions. Some applications for special means are also given.

Keywords—Hayashi's inequality, Hermite-Hadamard's inequality, Steffensen's inequality, Integral inequalities, Special means.

1. THE MAIN RESULT

To prove our main result, we shall need the following generalization of the well-known Steffensen's inequality which is due to Hayashi [1, pp. 311–312].

THEOREM 1. Let $f : [a, b] \rightarrow \mathbb{R}$ be a nonincreasing mapping on $[a, b]$ and $g : [a, b] \rightarrow \mathbb{R}$ an integrable mapping on $[a, b]$ with

$$0 \leq g(x) \leq A, \quad \text{for all } x \in [a, b].$$

Then, the following inequality holds

$$A \int_{b-\lambda}^b f(x) dx \leq \int_a^b f(x)g(x) dx \leq A \int_a^{a+\lambda} f(x) dx, \quad (1)$$

where

$$\lambda = \frac{1}{A} \int_a^b g(x) dx.$$

THEOREM 2. Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable mapping on I° and $[a, b] \subset I^\circ$ with $M = \sup_{x \in [a, b]} f'(x) < \infty$, $m = \inf_{x \in [a, b]} f'(x) > -\infty$ and $M > m$. If f' is integrable on $[a, b]$, then the following inequality holds

$$\left| \frac{1}{b-a} \int_a^b f(x) dx - \frac{f(a) + f(b)}{2} \right| \leq \frac{[f(b) - f(a) - m(b-a)][M(b-a) - f(b) + f(a)]}{2(M-m)(b-a)} \quad (2)$$

$$\leq \frac{(M-m)(b-a)}{8}. \quad (3)$$

PROOF. Let $f(x) = a - x$ and $g(x) = f'(x) - m$, and apply Hayashi's inequality, to obtain

$$(M-m) \int_{b-\lambda}^b (a-x) dx \leq I \leq (M-m) \int_a^{a+\lambda} (a-x) dx, \quad (4)$$

where

$$I = \int_a^b (a-x)(f'(x) - m) dx,$$

and

$$\lambda = \frac{1}{M-m} \int_a^b (f'(x) - m) dx = \frac{f(b) - f(a) - m(b-a)}{M-m}.$$

Since,

$$\int_{b-\lambda}^b (a-x) dx = \frac{1}{2} [(b-a-\lambda)^2 - (b-a)^2],$$

and

$$\int_a^{a+\lambda} (a-x) dx = -\frac{\lambda^2}{2},$$

the inequality (4) is the same as

$$\ell_1 \equiv (M-m) \left[\frac{(b-a-\lambda)^2 - (b-a)^2}{2} \right] \leq I \leq (M-m) \left[-\frac{\lambda^2}{2} \right] \equiv \ell_2.$$

Next, since

$$\begin{aligned} \frac{\ell_1 + \ell_2}{2} &= \frac{M-m}{2} \left[-\frac{\lambda^2}{2} + \frac{(b-a-\lambda)^2}{2} - \frac{(b-a)^2}{2} \right] \\ &= \frac{(M-m)[- \lambda(b-a)]}{2} = \frac{m(b-a)^2}{2} - \frac{(b-a)(f(b) - f(a))}{2}, \end{aligned}$$

and

$$I = \int_a^b f(x) dx - (b-a)f(b) + m \frac{(b-a)^2}{2},$$

it follows that

$$\begin{aligned} \left| I - \frac{\ell_1 + \ell_2}{2} \right| &= \left| \int_a^b f(x) dx - (b-a)f(b) + m \frac{(b-a)^2}{2} - m \frac{(b-a)^2}{2} + \frac{(b-a)(f(b) - f(a))}{2} \right| \\ &= \left| \int_a^b f(x) dx - (b-a) \frac{f(a) + f(b)}{2} \right|. \end{aligned} \quad (5)$$

We also have

$$\begin{aligned} \left| I - \frac{\ell_1 + \ell_2}{2} \right| &\leq \frac{\ell_2 - \ell_1}{2} \\ &= \frac{M-m}{2} \left[-\frac{\lambda^2}{2} + \frac{(b-a)^2}{2} - \frac{(b-a-\lambda)^2}{2} \right] \\ &= \frac{M-m}{2} [-\lambda^2 + (b-a)\lambda] \\ &= \frac{[f(b) - f(a) - m(b-a)][M(b-a) - f(b) + f(a)]}{2(M-m)}. \end{aligned} \quad (6)$$

Now on combining (5) and (6) we immediately obtain the inequality (2).

To prove (3), we define the mapping $g: \mathbb{R} \rightarrow \mathbb{R}$, $g(t) = -t^2 + (b-a)t$. It is clear that

$$g(t) \leq g\left(\frac{b-a}{2}\right) = \frac{(b-a)^2}{4}, \quad \text{for all } t \in \mathbb{R}. \quad (7)$$

We choose

$$t = \lambda = \frac{f(b) - f(a) - m(b-a)}{M-m},$$

so that

$$\frac{M-m}{2}g(\lambda) = \frac{[f(b) - f(a) - m(b-a)][M(b-a) - f(b) + f(a)]}{2(M-m)}. \quad (8)$$

From (7) and (8), we have

$$\frac{M-m}{2}g(\lambda) \leq \frac{(M-m)(b-a)^2}{8},$$

which in view of (2) proves the required inequality (3).

COROLLARY 3. *Let the conditions of Theorem 2 be satisfied, and $\|f'\|_\infty = \sup_{x \in [a,b]} |f'(x)| < \infty$. Then, the following inequality holds*

$$\begin{aligned} \left| \frac{1}{b-a} \int_a^b f(x) dx - \frac{f(a) + f(b)}{2} \right| &\leq \frac{\|f'\|_\infty^2 [(b-a)^2 - (f(b) - f(a))^2]}{4\|f'\|_\infty(b-a)} \\ &\leq \frac{\|f'\|_\infty(b-a)}{4}. \end{aligned}$$

PROOF. The corollary follows from Theorem 2 by choosing $m = -\|f'\|_\infty$ and $M = \|f'\|_\infty$.

REMARK 1. The first inequality in Corollary 3 was proved by Lyengar in 1938 (see [2, p. 471]) by means of geometrical arguments.

The case of convex mappings is embodied in the following corollary, which is very important in applications.

COROLLARY 4. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable convex mapping on I° and $[a, b] \subset I^\circ$ with $f'(b) \neq f'(a)$. Then, the following inequality holds*

$$\begin{aligned} 0 &\leq \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \\ &\leq \frac{[f(b) - f(a) - f'(a)(b-a)][f'(b)(b-a) - f(b) + f(a)]}{2(f'(b) - f'(a))(b-a)} \\ &\leq \frac{(f'(b) - f'(a))(b-a)}{8}. \end{aligned}$$

PROOF. The corollary follows from Theorem 2 and the observation that we can choose $m = f'(a)$ and $M = f'(b)$.

REMARK 2. For a convex mapping $f : [a, b] \rightarrow \mathbb{R}$, the double inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}$$

is well known in literature as Hermite-Hadamard's integral inequality [3] and references therein. The result of Corollary 3 gives a counterpart of the above inequality. Indeed, we have

$$\frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \geq 0,$$

when the mapping f is differentiable on I° , i.e., in the usual cases which are important in applications.

2. APPLICATIONS TO SPECIAL MEANS

For $\alpha, \beta > 0$ we recall the means

$$\begin{aligned}
 A(\alpha, \beta) &= \frac{\alpha + \beta}{2}, & \text{arithmetic mean,} \\
 G(\alpha, \beta) &= \sqrt{\alpha\beta}, & \text{geometric mean,} \\
 H(\alpha, \beta) &= \frac{2}{(1/\alpha) + (1/\beta)}, & \text{harmonic mean,} \\
 L(\alpha, \beta) &= \frac{\beta - \alpha}{\ln \beta - \ln \alpha}, & \text{logarithmic mean } (\alpha \neq \beta), \\
 I(\alpha, \beta) &= \frac{1}{e} \left(\frac{\beta\beta}{\alpha\alpha} \right)^{1/(\beta-\alpha)}, & \text{identric mean } (\alpha \neq \beta), \\
 L_p(\alpha, \beta) &= \left[\frac{\beta^{p+1} - \alpha^{p+1}}{(p+1)(\beta - \alpha)} \right]^{1/p}, & \text{generalized log-mean } (\alpha \neq \beta), p \neq -1, 0.
 \end{aligned}$$

Now we shall use the results of Section 1 to prove the following new facts for the above means.

PROPOSITION 1. *Let $p \geq 1$ and $0 \leq a \leq b$. Then, the following hold*

$$\begin{aligned}
 0 \leq A(a^p, b^p) - L_p^p(a, b) &\leq \frac{p}{2(p-1)} \frac{\left[L_{p-1}^{p-1}(a, b) - a^{p-1} \right] \left[b^{p-1} - L_{p-1}^{p-1}(a, b) \right]}{L_{p-2}^{p-2}(a, b)} \\
 &\leq \frac{p(p-1)}{8} (b-a)^2 L_{p-2}^{p-2}(a, b).
 \end{aligned}$$

PROOF. By Corollary 4 applied to the convex mapping $f(x) = x^p$, $p \geq 1$, we have

$$\begin{aligned}
 0 &\leq \frac{a^p + b^p}{2} - \frac{1}{b-a} \int_a^b x^p dx \\
 &\leq \frac{[b^p - a^p - pa^{p-1}(b-a)] [pb^{p-1}(b-a) - b^p + a^p]}{2p(b^{p-1} - a^{p-1})(b-a)} \\
 &\leq \frac{p(b^{p-1} - a^{p-1})(b-a)}{8}.
 \end{aligned}$$

Proposition 1 now follows from the facts that

$$b^p - a^p = p(b-a)L_{p-1}^{p-1}(a, b),$$

and

$$b^{p-1} - a^{p-1} = (p-1)(b-a)L_{p-2}^{p-2}(a, b).$$

PROPOSITION 2. *Let $0 < a < b$. Then, the following hold*

$$0 \leq L(a, b) - H(a, b) \leq L(a, b) \left(\frac{b-a}{a+b} \right)^2 \leq L(a, b) \frac{(b-a)^2}{4ab}. \quad (9)$$

PROOF. By Corollary 4 applied to the convex mapping $f(x) = 1/x$ on $[a, b] \subset (0, \infty)$, we have

$$\begin{aligned}
 0 &\leq \frac{1/a + 1/b}{2} - \frac{\ln b - \ln a}{b-a} \\
 &\leq \frac{[1/b - 1/a + ((b-a)/a^2)] [-(b-a)/b^2 - 1/b + 1/a]}{2(1/a^2 - 1/b^2)(b-a)} = R, \quad (\text{say}) \\
 &\leq \frac{(1/a^2 - 1/b^2)(b-a)}{8}.
 \end{aligned}$$

However, since

$$\frac{1}{b} - \frac{1}{a} + \frac{b-a}{a^2} = \frac{(b-a)^2}{a^2b},$$

and

$$-\frac{b-a}{b^2} - \frac{1}{b} + \frac{1}{a} = \frac{(b-a)^2}{ab^2},$$

it follows that

$$R = \frac{(b-a)^2}{2ab(a+b)}.$$

Consequently, we get

$$0 \leq H^{-1}(a, b) - L^{-1}(a, b) \leq \frac{(b-a)^2}{2ab(a+b)} \leq \frac{(b-a)^2(a+b)}{8a^2b^2},$$

which is equivalent to (9).

PROPOSITION 3. *Let $0 < a < b$. Then, the following holds*

$$0 \leq \ln I(a, b) - \ln G(a, b) \leq \frac{ab [\ln(a/b) + ((b-a)/a)] [\ln(b/a) - ((b-a)/b)]}{2(b-a)^2} \leq \frac{(b-a)^2}{8ab}.$$

PROOF. The proof follows from Corollary 4 applied to the convex mapping $f(x) = -\ln x$.

REFERENCES

1. D.S. Mitrinović, J.E. Pečarić and A.M. Fink, *Classical and New Inequalities in Analysis*, Kluwer, Dordrecht, (1993).
2. D.S. Mitrinović, J.E. Pečarić and A.M. Fink, *Inequalities Involving Functions and Their Integrals and Derivatives*, Kluwer, Dordrecht, (1991).
3. S.S. Dragomir, Two mappings in connection to Hadamard's inequality, *J. Math. Anal. Appl.* **167**, 49–56 (1992).